

Five Things Physicians and Patients Should Question

1

Don't perform unproven diagnostic tests, such as immunoglobulin G (IgG) testing or an indiscriminate battery of immunoglobulin E (IgE) tests, in the evaluation of allergy.

Appropriate diagnosis and treatment of allergies requires specific IgE testing (either skin or blood tests) based on the patient's clinical history. The use of other tests or methods to diagnose allergies is unproven and can lead to inappropriate diagnosis and treatment. Appropriate diagnosis and treatment is both cost effective and essential for optimal patient care.

2

Don't order sinus computed tomography (CT) or indiscriminately prescribe antibiotics for uncomplicated acute rhinosinusitis.

Viral infections cause the majority of acute rhinosinusitis and only 0.5 percent to 2 percent progress to bacterial infections. Most acute rhinosinusitis resolves without treatment in two weeks. Uncomplicated acute rhinosinusitis is generally diagnosed clinically and does not require a sinus CT scan or other imaging. Antibiotics are not recommended for patients with uncomplicated acute rhinosinusitis who have mild illness and assurance of follow-up. If a decision is made to treat, amoxicillin should be first-line antibiotic treatment for most acute rhinosinusitis.

3

Don't routinely do diagnostic testing in patients with chronic urticaria.

In the overwhelming majority of patients with chronic urticaria, a definite etiology is not identified. Limited laboratory testing may be warranted to exclude underlying causes. Targeted laboratory testing based on clinical suspicion is appropriate. Routine extensive testing is neither cost effective nor associated with improved clinical outcomes. Skin or serum-specific IgE testing for inhalants or foods is not indicated, unless there is a clear history implicating an allergen as a provoking or perpetuating factor for urticaria.

4

Don't recommend replacement immunoglobulin therapy for recurrent infections unless impaired antibody responses to vaccines are demonstrated.

Immunoglobulin (gammaglobulin) replacement is expensive and does not improve outcomes unless there is impairment of antigen-specific IgG antibody responses to vaccine immunizations or natural infections. Low levels of immunoglobulins (isotypes or subclasses), without impaired antigen-specific IgG antibody responses, do not indicate a need for immunoglobulin replacement therapy. Exceptions include IgG levels <150mg/dl and genetically defined/suspected disorders. Measurement of IgG subclasses is not routinely useful in determining the need for immunoglobulin therapy. Selective IgA deficiency is not an indication for administration of immunoglobulin.

5

Don't diagnose or manage asthma without spirometry.

Clinicians often rely solely upon symptoms when diagnosing and managing asthma, but these symptoms may be misleading and be from alternate causes. Therefore spirometry is essential to confirm the diagnosis in those patients who can perform this procedure. Recent guidelines highlight spirometry's value in stratifying disease severity and monitoring control. History and physical exam alone may over- or under-estimate asthma control. Beyond the increased costs of care, repercussions of misdiagnosing asthma include delaying a correct diagnosis and treatment.

Five More Things Physicians and Patients Should Question

6

Don't rely on antihistamines as first-line treatment in severe allergic reactions.

Epinephrine is the first-line treatment for anaphylaxis. Data indicate that antihistamines are overused as the first-line treatment of anaphylaxis. By definition, anaphylaxis has cardiovascular and respiratory manifestations, which require treatment with epinephrine. Overuse of antihistamines, which do not treat cardiovascular or respiratory manifestations of anaphylaxis, can delay the effective first-line treatment with epinephrine.

Epinephrine should be administered as soon as the diagnosis of anaphylaxis is suspected. Antihistamines are second-line supportive therapy for cutaneous non-life-threatening symptoms (hives), but do not replace epinephrine as the first-line treatment for anaphylaxis.

Fatalities during anaphylaxis have been associated with delayed administration of epinephrine.

7

Don't perform food IgE testing without a history consistent with potential IgE-mediated food allergy.

False or clinically irrelevant positive allergy tests for foods are frequent. Indiscriminate screening results in inappropriate avoidance of foods and wastes healthcare resources. IgE testing for specific foods must be driven by a history of signs or symptoms consistent with an IgE-mediated reaction after eating a particular food. Ordering IgE testing in individuals who do not have a history consistent with or suggestive for food allergy based on history frequently reveals positive tests that are unlikely to be clinically relevant. Testing, when done, should be limited to suspected foods.

The diagnostic utility of IgE testing for specific foods is optimal when a history compatible with or suggestive for the diagnosis of food allergy is present. In the absence of a compatible or suggestive history, the pre-test probability for a diagnosis of food allergy is low and a positive skin or in vitro IgE test does not establish a diagnosis of food allergy. Skin testing or serum testing for specific-IgE to food antigens has excellent sensitivity and high negative predictive value, but has low specificity and low positive predictive value.

Considering that 50 to 90 percent of presumed cases of food allergy do not reflect IgE-mediated (allergic) pathogenesis and may instead reflect food intolerance or symptoms not causally associated with food consumption, ordering panels of food tests leads to many incorrectly identified food allergies and inappropriate recommendations to avoid foods that are positive on testing.

8

Don't routinely order low- or iso-osmolar radiocontrast media or pretreat with corticosteroids and antihistamines for patients with a history of seafood allergy, who require radiocontrast media.

Although the exact mechanism for contrast media reactions is unknown, there is no cause and effect connection with seafood allergy. Consequently there is no reason to use more expensive agents or pre-medication before using contrast media in patients with a history of seafood allergy. A prior history of anaphylaxis to contrast media is an indication to use low- or iso-osmolar agents and pretreat with corticosteroids and antihistamines.

Patients with a history of seafood allergy are not at elevated risk for anaphylaxis from iodinated contrast media. Similarly, patients who have had anaphylaxis from contrast media should not be told that they are allergic to seafood.

Patients with a history of seafood allergy who are labeled as being at greater risk for adverse reaction from contrast infusions experience considerable morbidity from unnecessary precautions – including but not limited to denying them indicated roentgenographic procedures and adverse effects from pretreatment with antihistamine and/or corticosteroid medications.

Regardless of whether these patients truly have IgE-mediated allergies to seafood (crustacean), there is no evidence in the medical literature that indicates they are at elevated risk for anaphylaxis from contrast infusion compared with the history-negative general population.

In a random telephone survey of 5,529 households with a census of 14,948 individuals, seafood allergy was reported by 3.3 percent of survey respondents. According to current U.S. population estimates for 2013, this corresponds to 10,395,000 Americans.

The mechanism for anaphylaxis to radio-iodinated contrast media relates to the physiochemical properties of these media and is unrelated to its iodine content. Further, although delayed-type hypersensitivity (allergic contact dermatitis) reactions to iodine have rarely been reported, IgE-mediated reactions to iodine have not, and neither type of reaction would be related to IgE-mediated shellfish allergy nor to contrast media reactions. Patients with a history of prior anaphylaxis to contrast media are at elevated risk for anaphylactic reaction with re-exposure to contrast media.

Patients with asthma or cardiovascular disease, or who are taking beta blockers, are at increased risk for serious anaphylaxis from radiographic contrast media.

Don't routinely avoid influenza vaccination in egg-allergic patients.

Of the vaccines that may contain egg protein (measles, mumps, rabies, influenza and yellow fever), measles, mumps and rabies vaccines have at most negligible egg protein; consequently no special precautions need to be followed in egg-allergic patients for these vaccines. Studies in egg-allergic patients receiving egg-based inactivated influenza vaccine have not reported reactions; consequently egg-allergic patients should be given either egg-free influenza vaccine or should receive egg-based influenza vaccine with a 30-minute post-vaccine observation period. Egg-allergic patients receiving the yellow fever vaccine should be skin tested with the vaccine and receive the vaccine with a 30-minute observation period if the skin test is negative. If positive, the vaccine may be given in graded doses with appropriate medical observation.

Egg protein is present in influenza and yellow fever vaccines and in theory could cause reactions in egg-allergic patients. However, in 27 published studies collectively 4,172 patients with egg allergy received 4,729 doses of egg-based inactivated influenza vaccine (IIV) with no cases of anaphylaxis, including 513 with severe egg allergy who uneventfully received 597 doses. The CDC's Advisory Committee on Immunization Practices recommends that egg-allergic persons receive IIV as a single dose without prior vaccine skin testing and be observed for 30 minutes afterwards for any possible allergic reaction. If the reaction to the ingestion of eggs was hives only, the vaccine can be administered in a primary care setting, whereas if the reaction to the ingestion of eggs was more severe, the vaccine should be administered in an allergist/immunologist's office. Two new IIVs not grown in eggs have been approved for patients 18 years and older: Flucelvax, prepared from virus propagated in cell culture, and Flublok, recombinant hemagglutinin proteins produced in an insect cell line. For egg-allergic patients 18 years of age and older, either egg-based IIV can be used with the precautions above or egg-free IIV can be used.

Measles and mumps vaccines (and Purified Chick Embryo Cell [PCEC] rabies vaccine) are grown in chick embryo fibroblast cultures and contain negligible or no egg protein. Thus, MMR and PCEC rabies vaccine can be administered to egg-allergic recipients in the usual manner.

Per the Yellow Fever vaccine package insert, egg-allergic recipients should be skin tested with the vaccine prior to administration. If negative, the vaccine can be given in the usual manner, but the patient should be observed for 30 minutes afterward. If the vaccine skin test is positive, the vaccine can be given in graded doses under appropriate medical observation.

Don't overuse non-beta lactam antibiotics in patients with a history of penicillin allergy, without an appropriate evaluation.

While about 10 percent of the population reports a history of penicillin allergy, studies show that 90 percent or more of these patients are not allergic to penicillins and are able to take these antibiotics safely. The main reason for this observation is that penicillin allergy is often misdiagnosed and when present wanes over time in most (but not all) individuals. Patients labeled penicillin-allergic are more likely to be treated with alternative antibiotics (such as vancomycin and quinolones), have higher medical costs, experience longer hospital stays, and are more likely to develop complications such as infections with vancomycin-resistant enterococcus (VRE) and *Clostridium difficile*.

Evaluation for specific IgE to penicillin can be carried out by skin testing. Ideally, penicillin skin testing should be performed with both major and minor determinants. The negative predictive value of penicillin skin testing for immediate reactions approaches 100 percent, whereas the positive predictive value is between 40 and 100 percent. The usefulness of in vitro tests for penicillin-specific IgE is limited by their uncertain predictive value. They are not suitable substitutes for penicillin skin testing.

By identifying the overwhelming majority of individuals who can safely receive penicillin and penicillin-like drugs, we can improve the appropriateness of antibiotic therapy and clinical care outcomes.

How This List Was Created

The American Academy of Allergy, Asthma & Immunology (AAAAI) Executive Committee created a task force to lead work on Choosing Wisely consisting of board members, the AAAAI President and Secretary/Treasurer and AAAAI participants in the Joint Task Force on Practice Parameters. Through multiple society publications and notifications, AAAAI members were invited to offer feedback and recommend elements to be included in the list. A targeted email was also sent to an extended group of AAAAI leadership inviting them to participate.

The work group reviewed the submissions to ensure the best science in the specialty was included. Based on this additional members were recruited for their expertise. Suggested elements were considered for appropriateness, relevance to the core of the specialty, potential overuse of resources and opportunities to improve patient care. They were further refined to maximize impact and eliminate overlap, and then ranked in order of potential importance both for the specialty and for the public. Finally, the work group chose its top five recommendations which were then approved by the Executive Committee. AAAAI's disclosure and conflict of interest policy can be found at www.aaaai.org.

Sources

- 1 Cox L, Williams PB, Sicherer S, et al. Pearls and pitfalls of allergy diagnostic testing: report from the American College of Allergy, Asthma and Immunology/ American Academy of Allergy, Asthma & Immunology Specific IgE Test Task Force. *Ann All Asthma Immunol*. 2008;101:580–92
Bernstein I, Li J, Bernstein D et al. Allergy diagnostic testing: an updated practice parameter. *Ann All Asthma Immunol* 2008;100:s1–148.
Terr AI. Unconventional theories and unproven methods in allergy. In: *Allergy Principles and Practice*, 7th Ed, 97:1691–1709.
- 2 Ahovuo-Saloranta A, Borisenko OV, Kovanen N, et al. Antibiotics for acute maxillary sinusitis. *Cochrane database of systematic reviews* 2008:CD000243.
American College of Radiology ACR Appropriateness Criteria® for Sinonasal Disease, 2009 http://www.acr.org/SecondaryMainMenuCategories/quality_safety/app_criteria/pdf/ExpertPanelonNeurologicImaging/SinonasalDisease.aspx; 2009.
- 3 Wanderer, AA, Bernstein, IL, Goodman, DL, et al. The Diagnosis and Management of Urticaria: a Practice Parameter. *Ann Allergy Asthma Immunol* 2000;85:521–44.
Tarbox JA, Gutta RC, Ching EL, Radojicic C, Lang DM. Utility of routine laboratory testing in management of chronic urticaria/angioedema. *Ann Allergy Asthma Immunol* 2011, 107: 239–43.
Bernstein IL, Li, JT, Bernstein DI et al. Allergy diagnostic testing: an updated practice parameter. *Ann Allergy Asthma Immunol*. 2008 Mar;100(3 Suppl 3):S1–148.
Kozel MM, Bossuyt PM, Mekkes JR, Bos JD. Laboratory tests and identified diagnoses in patients with physical and chronic urticaria and angioedema: A systematic review. *J Am Acad Dermatol*. 2003 Mar;48(3):409–16.
- 4 Orange, JS et al. Use of intravenous immunoglobulin in human disease: a review of evidence by members of the Primary Immunodeficiency Committee of the American Academy of Allergy, Asthma and Immunology. *JACI* 117:S525–S553, 2006.
Ballow, M. “Immunoglobulin Therapy: Replacement and Immunomodulation” in *Clinical Immunology*, Third Edition Rich RR (Editor), Chapter 85, pp. 1265–1280, 2008.
Stiehm ER, Orange JS, Ballow M, Lehman H. Therapeutic use of immunoglobulins. *Adv Pediatr* 2010;57:185–218.
Bonilla FA, Bernstein IL, Khan DA, Ballas ZK, Chinen J, Frank MM, et al. Practice parameter for the diagnosis and management of primary immunodeficiency. *Annals of Allergy, Asthma & Immunology*. 2005;94 (Suppl 1):S1–S63.
- 5 National Asthma Education and Prevention Expert Panel Report 3: Guidelines for the diagnosis and Management of Asthma. NIH Publication Number 08–5846 October 2007.
Li J, Oppenheimer J, Bernstein IL et al. Attaining asthma control. A practice parameter. *J Allergy Clin Immunol*. 2005;115:S3–11.
Global strategy for asthma management and prevention: GINA executive summary *Eur Respir J* 2008 31:143–178.
Fuhlbrigge A, Kitch B, Paltelet D et al. FEV1 is associated with risk of asthma attacks in a pediatric population. *J Allergy Clin Immunol*. 2001;107:61–6.
Magadle R The Risk of Hospitalization and Near-Fatal and Fatal Asthma in Relation to the Perception of Dyspnea *Chest*. 2002;121:329–333.
- 6 Lieberman P, Nicklas RA, Oppenheimer J, Kemp SF, Lang DM, Bernstein DI, Bernstein JA, Burks AW, Feldweg AM, Fink JN, Greenberger PA, Golden DB, James JM, Kemp SF, Ledford DK, Lieberman P, Sheffer AL, Bernstein DI, Blessing-Moore J, Cox L, Khan DA, Lang D, Nicklas RA, Oppenheimer J, Portnoy JM, Randolph C, Schuller DE, Spector SL, Tilles S, Wallace D. The diagnosis and management of anaphylaxis practice parameter 2010 update. *J Allergy Clin Immunol*. 2010 Sep;126(3):477-80.e1–42.
Sampson HA, Muñoz-Furlong A, Campbell RL, Adkinson NF Jr, Bock SA, Branum A, Brown SG, Camargo CA Jr, Cydulka R, Galli SJ, Gidudu J, Gruchalla RS, Harlor AD Jr, Hepner DL, Lewis LM, Lieberman PL, Metcalfe DD, O'Connor R, Muraro A, Rudman A, Schmitt C, Scherrer D, Simons FE, Thomas S, Wood JP, Decker WW. Second symposium on the definition and management of anaphylaxis: summary report – Second National Institute of Allergy and Infectious Diseases/ Food Allergy and Anaphylaxis Network symposium. *J Allergy Clin Immunol*. 2006 Feb;117(2):391–7.
Kemp SF, Lockey RF, Simons FE; World Allergy Organization ad hoc Committee on Epinephrine in Anaphylaxis. Epinephrine the drug of choice for anaphylaxis. A statement of the World Allergy Organization. *Allergy*. 2008 Aug;63(8):1061–70.
Cox L, Nelson H, Lockey R, Calabria C, Chacko T, Finegold I, Nelson M, Weber R, Bernstein DI, Blessing-Moore J, Khan DA, Lang DM, Nicklas RA, Oppenheimer J, Portnoy JM, Randolph C, Schuller DE, Spector SL, Tilles S, Wallace D. Allergen immunotherapy: a practice parameter third update. *J Allergy Clin Immunol*. 2011 Jan;127(1 Suppl):s1–55.
Golden DB, Moffitt J, Nicklas RA, Freeman T, Graft DF, Reisman RE, Tracy JM, Bernstein D, Blessing-Moore J, Cox L, Khan DA, Lang DM, Oppenheimer J, Portnoy JM, Randolph C, Schuller DE, Spector SL, Tilles SA, Wallace D; Joint Task Force on Practice Parameters; American Academy of Allergy, Asthma & Immunology (AAAAI); American College of Allergy, Asthma & Immunology (ACAAI); Joint Council of Allergy, Asthma and Immunology. Stinging insect hypersensitivity: a practice parameter update 2011. *J Allergy Clin Immunol*. 2011 Apr; 127(4):852–4.
Clark S, Long AA, Gaeta TJ, Camargo CC. Multicenter study of emergency department visits for insect sting allergies. *J Allergy Clin Immunol*. 2005;116:643–9.

7

Bernstein IL, Li JT, Bernstein DI, Hamilton R, Spector SL, Tan R, Sicherer S, Golden DB, Khan DA, Nicklas RA, Portnoy JM, Blessing-Moore J, Cox L, Lang DM, Oppenheimer J, Randolph CC, Schuller DE, Tilles SA, Wallace DV, Levetin E, Weber R; American Academy of Allergy, Asthma and Immunology; American College of Allergy, Asthma and Immunology. Allergy diagnostic testing: an updated practice parameter. *Ann Allergy Asthma Immunol.* 2008 Mar;100(3 Suppl 3):S1–148.

NIAID-Sponsored Expert Panel, Boyce JA, Assa'ad A, Burks AW, Jones SM, Sampson HA, Wood RA, Plaut M, Cooper SF, Fenton MJ, Arshad SH, Bahna SL, Beck LA, Byrd-Bredbenner C, Camargo CA Jr, Eichenfield L, Furuta GT, Hanifin JM, Jones C, Kraft M, Levy BD, Lieberman P, Luccioli S, McCall KM, Schneider LC, Simon RA, Simons FE, Teach SJ, Yawn BP, Schwanger JM. Guidelines for the diagnosis and management of food allergy in the United States: report of the NIAID-sponsored expert panel. *J Allergy Clin Immunol.* 2010 Dec;126 (6 Suppl):S1–58.

8

American Academy of Asthma, Allergy and Immunology. Food allergy: a practice parameter. *Ann Allergy Asthma Immunol.* 2006 Mar;96:S1–68.

Lieberman P, Nicklas RA, Oppenheimer J, Kemp SF, Lang DM. The diagnosis and management of anaphylaxis practice parameter: 2010 update. *J Allergy Clin Immunol.* 2010 Aug 21;126(3):477–522.

Solensky R, Khan DA. Drug allergy: an updated parameter. *Ann Allergy Asthma Immunol.* 2010 Oct;105(4):259–73.

Sicherer S, Munoz-Furlong A, Sampson H. Prevalence of seafood allergy in the United States determined by a random telephone survey. *J Allergy Clin Immunol.* 2004;114:159–65.

Greenberger P. Prophylaxis against repeated radio contrast media reaction in 857 cases. *Arch Intern Med.* 1985;145:2197–200.

Sicherer SH. Risk of severe allergic reactions from the use of potassium iodide for radiation emergencies. *J Allergy Clin Immunol.* 2004;114:1395–7.

Lang DM, Alpern MB, Visintainer PF, Smith ST. Elevated risk for anaphylactoid reaction from radiographic contrast media associated with both beta blocker exposure and cardiovascular disorders. *Arch Intern Med.* 1993;153:2033–40.

9

Des Roches A, Paradis L, Gagnon R, Lemire C, Bégin P, Carr S, Chan ES, Paradis J, Frenette L, Ouakki M, Benoît M, De Serres G; PCIRN (Public Health Agency of Canada/Canadian Institutes of Health Research Influenza Research Network). Egg-allergic patients can be safely vaccinated against influenza. *J Allergy Clin Immunol.* 2012 Nov;130(5):1213–1216.

Centers for Disease Control and Prevention (CDC). Prevention and control of influenza with vaccines: recommendations of the Advisory Committee on Immunization Practices (ACIP)—United States, 2012–13 influenza season. *MMWR Morb Mortal Wkly Rep.* 2012 Aug 17;61(32):613–8.

FLUCELVAX (Novartis) Package Insert. 2012.

FLUBLOK (Protein Sciences) Package Insert. 2013.

American Academy of Pediatrics. Red Book: 2012 report of the Committee on Infectious Diseases. Pickering LK, ed. 29th ed. Elk Grove Village, IL: American Academy of Pediatrics; 2012. 936 p.

Kelso JM, Greenhawt MJ, Li JT, Nicklas RA, Bernstein DI, Blessing-Moore J, Cox L, Khan D, Lang DM, Oppenheimer J, Portnoy JM, Randolph CR, Schuller DE, Spector SL, Tilles SA, Wallace D. Adverse reactions to vaccines practice parameter 2012 update. *J Allergy Clin Immunol.* 2012 Jul;130(1):25–43.

YF-VAX (Sanofi Pasteur) Package Insert. 2010.

10

Solensky R, Khan DA. Drug allergy: an updated parameter. *Ann Allergy Asthma Immunol.* 2010 Oct;105(4):259–73.

Solensky R. Penicillin allergy as a public health measure. *J Allergy Clin Immunol.* 2013 Dec 8. pii:S0091-6749(13)01646–1.

Macy E, Contreras R. Healthcare utilization and serious infection prevalence associated with penicillin “allergy” in hospitalized patients: a cohort study. *J Allergy Clin Immunol.* 2013 Nov 1. pii:S0091–6749(13)01467–X.

Park MA, Markus PJ, Matesic D, Li JTC. Safety and effectiveness of a preoperative allergy clinic in decreasing vancomycin use in patients with a history of penicillin allergy. *Ann Allergy Asthma Immunol.* 2006;97:681–7.

About the ABIM Foundation

The mission of the ABIM Foundation is to advance medical professionalism to improve the health care system. We achieve this by collaborating with physicians and physician leaders, medical trainees, health care delivery systems, payers, policymakers, consumer organizations and patients to foster a shared understanding of professionalism and how they can adopt the tenets of professionalism in practice.

To learn more about the ABIM Foundation, visit www.abimfoundation.org.



About the American Academy of Asthma, Allergy and Immunology

The American Academy of Allergy, Asthma & Immunology (AAAAI) represents allergists, asthma specialists, clinical immunologists, allied health professionals, and others with a special interest in the research and treatment of allergic and immunologic diseases. Established in 1943, the AAAAI has more than 6,500 members in the United States, Canada, and 60 other countries.

For more information or questions, please visit www.aaaai.org.

